

Australian and New Zealand
Journal of Psychiatry

informa
healthcare

**Psychiatric Symptoms and Dissociation in Conversion,
Somatization and Dissociative Disorders**

Journal:	<i>Australian and New Zealand Journal of Psychiatry</i>
Manuscript ID:	ANP-2008-00241.R1
Manuscript Type:	Regular Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Espirito Santo, Helena; Instituto Superior Miguel Torga, Psicologia; Universidade do Porto, Instituto de Ciências Biomédicas Abel Salazar Abreu, Jose Luis; Universidade de Coimbra, Faculdade de Medicina
Keywords:	Conversion disorder, Dissociative disorder, Dissociation, Somatization disorder, Hysteria

 **scholarONE™**
Manuscript Central

Psychiatric Symptoms and Dissociation in Conversion, Somatization and Dissociative
Disorders

Helena Espirito-Santo¹, PsyD., M. D.
José Luís Pio-Abreu², MD., Phd., Professor

Department of Psychology¹
Instituto Superior Miguel Torga
R. Augusta, nº 46
Coimbra, Portugal
And
Instituto de Ciências Biomédicas de Abel Salazar,
Universidade do Porto, Portugal

Hospitais da Universidade de Coimbra ²
Faculdade de Medicina,
Universidade de Coimbra, Portugal

For submission to: Australian and New Zealand Journal of Psychiatry
Date: 28 outubro 2008

Corresponding Author: Doutora Helena Espirito Santo
Departamento de Psicologia
Instituto Superior Miguel Torga, Coimbra
Largo da Cruz de Celas, 1
3000-132 Coimbra
Portugal
Tel: (+351) 239 488 030
Fax: (+351) 239 488 031
Email: helenum@gmail.com

Abstract

Objective: Conversion, dissociation and somatization are historically related in the long established concept of hysteria. Somewhere along the way they were separated due to the Cartesian dualistic view. Our aim was to compare these pathologies and investigate whether symptoms of these pathologies overlap in their clinical appearance in a Portuguese sample.

Method: Twenty-six patients with conversion disorder, 38 with dissociative disorders, 40 with somatization disorder, and a comparison group of 46 patients having other psychiatric disorders answered questions about dissociation (Dissociative Experiences Scale), somatoform dissociation (Somatoform Dissociation Questionnaire), and psychopathological symptoms (Brief Symptom Inventory).

Results: Dissociative and somatoform symptoms were significantly more frequent in dissociative and conversion disorder than in somatization disorder and controls. There were no significant differences between dissociative and conversion patients.

Conclusions: Conversion disorder is closely related to dissociative disorders. These results support the ICD-10 categorisation of conversion disorder among dissociative disorders and the hypothesis of analogous psychopathological processes in conversion and dissociative disorders versus somatization disorder.

Keywords Conversion disorder; Dissociative disorder; Somatization disorder, Dissociation; Hysteria

Introduction

Hysteria has been an important and well-accepted disease for 3900 years, with an extraordinary and irregular conceptual evolution that ended in its disappearance from the majority of scientific writings [1]. In 1980, hysteria was replaced by distinct illnesses in the DSM and ICD, namely somatization, dissociative, and conversion. The DSM-IV-TR [2] classifies conversion disorder within somatoform disorders, along with somatization disorder. In the ICD-10 conversion disorder is categorised as a dissociative disorder (keeping its “hysteria” designation), and somatization disorder goes with somatoform disorders [3]. However, many investigators noted the great overlap between dissociative and somatoform disorders, as defined in DSM and ICD [4-8]. Several investigations showed that dissociative disorders may present with somatic symptoms [7, 9-17] and conversion symptoms [7, 10, 11, 18]. Conversely, many patients with somatization disorder also have dissociative symptoms [17, 19], especially if they have been abused [17, 20]. Additionally, many patients with conversion disorder report dissociative symptoms [8, 21-23], namely patients with pseudoneurological conversion conditions [24-29]. There is a great overlap between these disorders, and many studies have defended the inclusion of the conversion disorder in the group of dissociative disorders [6, 30, 31], like Pierre Janet’s perspective [32]; while others have supported the differentiation between conversion and somatization [33].

Dissociation, somatic and psychological, seems to be the underlying mechanism of these three pathologies. The mechanism of *psychoform* dissociation involves the loss of the integration of consciousness, memory, identity, and perception of the environment [2] (American Psychiatric Association, 1994). *Somatoform* dissociation is viewed as the lack

of integration of somatic experiences, functions and responses [34-38]. The most extensively employed self-report measures of dissociation, the Dissociative Experiences Scale (DES) [39] and Somatoform Dissociation Questionnaire (SDQ-20) [36], were developed to measure psychoform and somatoform dissociation respectively.

As far as we know, no other study has investigated the three hysterical nosological presentations, nor compared them with other psychopathological disorders in terms of somatoform and psychoform dissociation. This study was undertaken to assess the common and different features of somatization, dissociative and conversion disorders, studying the phenomena of psychoform dissociation, somatoform dissociation, and general psychopathology, and to compare them with other psychiatric conditions.

Material and Methods

We present the cross-sectional and self-report data collected.

Subjects

We selected 151 patients from amongst 1162 consecutive cases registered between 2005 and 2006 from three mental health centres; they met DSM-IV criteria for conversion disorder (n=26), dissociative disorders (n=39), somatization disorder (n=40) and other psychiatric disorders (n=46). Twenty-seven were inpatients and 123 were outpatients. We ruled out patients with psychotic disorders, substance abuse disorders, bipolar disorder, personality disorders, and those under 18 years old. Expert clinicians with several years of training (mean practice of 19.7 years) performed a longitudinal evaluation using all data available (LEAD methodology) [40]. Because of logistical factors, only the diagnoses of 62 of these patients were confirmed with a Portuguese version of the

Dissociative Disorders Interview Schedule (DDIS)[41]. Due to discharge, two patients did not fill out the SDQ-20. None of the other psychiatric patients met criteria for any of the above disorders. All patients gave written informed consent, and anonymity was preserved, according to the Code of Medical Ethics of the World Medical Association Declaration of Helsinki.

The demographic characteristics of the samples can be seen in Table 1. There were no significant differences in ages among the four groups ($F=1.81$, $df=3$, $p>0.05$), nor between genders ($\chi^2=1.14$, $df=3$, $p>0.05$), marital status ($\chi^2=3.10$, $df=3$, $p>0.05$), school years ($F=1.82$, $df=3$, $p>0.05$), or inpatients/outpatients ($\chi^2=7.07$, $df=3$, $p>0.05$). Table 2 shows the sub sample diagnosis frequencies.

Instruments

The Dissociative Experiences scale: Portuguese version. The DES is a self-administered 28-item questionnaire to measure the frequency of dissociative experiences [39], with subjects answering by circling the percentage of time they experience dissociation (from 0 to 100 in increments of 10%). The Portuguese version has good reliability (Cronbach’s $\alpha=0.94$) [42].

The *Somatoform Dissociation Questionnaire (SDQ-20)* is a 20-item self-administered tool to evaluate the intensity of somatoform dissociation [36]. Reliability of the Portuguese form was 0.88 (Cronbach’s α) [43].

The *Brief Symptom Inventory (BSI)* [44] is a 53-item self-report clinical rating scale that measures psychological distress (Global Severity Index) and nine dimensions: somatization, interpersonal sensitivity, anxiety, phobic anxiety, obsessive-compulsive symptoms, depression, hostility paranoid ideation, and psychoticism. We used the

Portuguese threshold [45] of 1.7 or greater for each subscale as an indication of symptom severity. These symptom subscales do not correspond to psychiatric diagnosis.

Cronbach's alpha for the subscales ranged from 0.62 (phobic anxiety) to 0.80 (somatization).

The *Dissociative Disorders Interview Schedule* (DDIS) [41] is a 131-item semi-structured interview that identifies all dissociative disorders, somatization disorder, and conversion disorder according to DSM-IV diagnoses, and a Portuguese adaptation (sensitivity rate 84%, specificity 100%, and overall kappa 0.83) [46].

Statistical Analysis

Data were analysed with Statistical Package for the Social Sciences (SPSS 11.0.3, for Mac OS X). Subjects with the four diagnoses were compared with each other according to the frequency (mean DES scores) and severity (above cut-off score 30) of dissociative experiences; mean SDQ-20 scores, and severity; distress and psychopathological symptoms from the BSI. None of these measures were normally distributed, therefore Kruskal-Wallis tests were used, followed by post-hoc Mann-Whitney tests on each pair of groups and adjusted *p* value with Bonferroni method (*p* value divided by the six paired comparisons made; in this way a significant level was set at $p < 0.008$).

Results

Results are described through the DES, SDQ and BSI measures used in this study.

Psychoform dissociation – DES

The DES ratings of all 151 patients ranged from 1.92 to 72.69, with a mean±SD of 26.82; there were no significant differences ($U=23.42$, $p=0.935$) between the mean DES scores of women (26.32 ± 14.03) and men (27.97 ± 17.55) or between single (26.23 ± 15.34) and married patients (27.69 ± 14.90) ($U=2555.50$, $p=0.579$). Inpatients (mean±SD= 30.06 ± 15.40) and outpatients (mean±SD= 26.10 ± 15.06) did not differ significantly ($U=1425.00$, $p=0.249$). There was no significant correlation between age and the scale scores ($\rho=0.138$, $p=0.094$) or between number of school years and scale scores ($\rho=0.102$, $p=0.214$). Thirty-eight percent of the patients got a score of 30 or above. There were no significant differences between higher and lower scores in terms of demographic characteristics.

Mean±SD DES scores were higher for the conversion patients, followed by dissociative disorders, somatization, and the comparison group. These scores differed significantly overall ($H=68.86$, $df=3$, $p<0.001$, Kruskal-Wallis one-way analysis of variance by ranks). Pairwise post-hoc U tests showed that dissociative and conversion patients did not differ, nor did somatizing patients from the comparison group. The other group comparisons, however, were significantly different. These results are shown in Table 3.

The proportions of patients from the diagnostic categories with above cut-off scores on the DES are presented in Table 4. Conversion and dissociative patients had significantly ($H=58.73$, $df=3$, $p<0.001$) more severe dissociative experiences than patients with somatization or other psychiatric disorders.

Somatoform dissociation-SDQ

The SDQ ranged from 20.00 to 76.00 and had a mean±SD of 32.00 ± 11.29 for all patients. Women scored 34.32 ± 11.73 and men 33.56 ± 10.30 and their differences were not

significant ($U=2145.00$, $p=0.394$); single patients scored 33.01 ± 9.58 and married 35.67 ± 13.34 , and their difference was also non-significant ($U=2453.00$, $p=0.465$). Inpatients had significantly higher ($U=1215.00$, $p=0.038$) somatoform dissociation scores ($\text{mean}\pm\text{SD}=39.15\pm14.21$) than outpatients ($\text{mean}\pm\text{SD}=32.96\pm10.26$). There was no significant correlation between scale scores and age ($r=0.121$, $p=0.142$). However, less educated subjects had higher scores ($r=-0.231$, $p=0.005$). A score of 35 or above was attained by 38% of the patients.

Conversion patients also had higher mean \pm SD SDQ scores, followed by dissociatives, then somatizing, and then other psychiatric patients. These scores differed significantly ($H=20.46$, $df=3$, $p<0.001$). The post-hoc Mann-Whitney tests revealed significant group differences between dissociative and somatizing, dissociative and controls, and between conversion and controls. Once again, dissociative and conversion patients did not differ from each other, nor did somatizing from controls. Conversion patients did not differ from somatizing participants (see Table 3).

More conversion patients scored higher than 35, followed by dissociative, then somatization and other psychiatric patients; these differences were significant ($H=20.14$, $df=3$, $p<0.001$) (see Table 4).

Psychopathology

The BSI data is presented in Table 3. The mean GSI scores were below the 1.7 threshold for the four groups. Conversion patients scored higher than 1.7 in anxiety and obsessive-compulsive symptoms. Dissociative patients had higher levels in obsessive, depression and paranoid symptoms. Somatizing patients scored higher in somatization and obsessive

symptoms. As a group, patients with other psychiatric disorders did not score high in any subscales. There were no significant differences between any paired groups in mean±SD GSI scores. We found significant differences only in somatization symptoms, with conversion patients and somatization patients revealing higher levels than patients with other psychiatric diagnoses.

Discussion

The aim of this study has been to analyze the relationship between conversion disorder, dissociative disorders and somatization disorder across the spectrums of psychoform dissociation, somatoform dissociation, and general psychopathology.

As far as we know, there is only one investigation that has compared conversion and somatization disorder with regard to psychoform dissociative symptoms, and reported no significant differences [21]; our results do not support this investigation: the patients with conversion had higher scores.

We found no studies that compared dissociative and somatization disorder, considering psychoform dissociation. Our findings reveal that dissociative patients scored higher than the somatization subjects.

We also found that patients with conversion and dissociative disorder patients differed significantly from somatization ones regarding somatoform dissociative symptoms; no other study measured this aspect.

Our main finding was a closer association between conversion and dissociative disorders, than between conversion and somatization disorder. This result is supported by: (1) greater psychoform and somatoform dissociation in conversion and dissociation; (2) a

lack of differences between conversion and dissociative disorders using measures of psychoform dissociation, somatoform dissociation and general psychopathology; (3) differences between conversion and somatization disorder in psychoform but not somatoform dissociation. Additionally, there is a common element to the three groups of pathology, and this is the obsessive symptoms.

These are timely findings, considering the discussion regarding somatoform disorder classification. There are both practical [47, 48] and theoretical issues [49]. Some argue that conversion is misplaced in DSM-IV [6, 7, 8, 37] and should be reunited with dissociative disorders [6, 7, 22, 50-52] or that it definitely has a dissociative component [21, 23, 53]. We also propose that conversion is re-conceptualised as a particular type of dissociative disorder, with a somatization aspect. We suggest that psychoform and somatoform dissociation are specific mechanisms of dissociative and conversion disorders.

There were some limitations to our study. Given the small sample size, it is possible that some statistically significant differences occurred by chance, thus our conclusions are limited to general trends, and larger replication studies are required.

Also, our demographics are disproportionate in relation to the population; despite the fact that we had no significant differences in the proportions of demographic characteristics across the four groups, this selection bias may limit the generalisation of the study.

However, we know that women are more likely to search for medical or psychological help [54] (WHO, 2006), and are usually over-represented in these studies. Moreover, men and women had no significant differences in mean DES or SDQ scores or in the proportion of higher scores. We are aware that younger subjects tend to report higher

1
2
3 levels of psychological dissociation [34, 55-59] and somatoform dissociation [16, 60].
4
5
6 The same tendency occurs in less educated subjects for somatoform dissociation [16, 60].
7
8 These possible selection biases were controlled: there was no correlation between age or
9
10 number of school years and scores of the DES or SDQ, except for the proportion of
11
12 patients with an elevated SDQ score, who were less educated. Thus, it is important to
13
14 control the level of education in further studies. Some studies show a relationship
15
16 between being single and psychological dissociation [57-59], and somatoform
17
18 dissociation [57]. Again, there were no significant differences in mean DES or SDQ
19
20 scores or in the proportion of higher scores between single and married subjects.
21
22
23 A potential bias may have occurred from the self-report measures. It is possible that
24
25 symptom exaggeration and contamination with social desirability took place, since we
26
27 only used the structured interview with 62 subjects (41%). In addition, we recognize that
28
29 individuals with psychiatric conditions may have a co-morbid dissociative disorder which
30
31 goes undiagnosed unless a structured interview takes place [61]. Moreover, we used wide
32
33 psychiatric categories instead of specific diagnosis, which could introduce unmeasured
34
35 confusion. In fact, it has been demonstrated that conversion disorder could be a
36
37 heterogeneous diagnostic category [62]. Additionally, the evaluators were not blind to the
38
39 diagnosis. Therefore, a growing area for replication would be to use structured interviews
40
41 combined with self-report measures, do more research on specific diseases, and involve
42
43 blinded interviewers, unaware of the investigation hypothesis.
44
45
46 Despite these limitations, there are also strengths in this study. The participants were
47
48 inpatients and outpatients: knowing that inpatients are more likely to have psychiatric co-
49
50 morbidity, outpatients could have brought fewer biases.
51
52
53
54
55
56
57
58
59
60

We used, as well, a consecutive sampling procedure, which made the sample closer to a community sample.

Finally, this study was the first to explore the three types of hysteria through psychological and somatoform dissociation.

Acknowledgements: None

For Review Only

References

1. Veith I. *Hysteria, the history of a disease*. Chicago: The University of Chicago Press, 1965.

2. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 4th ed., text rev. Washington, D.C: APA, 2000.

3. World Health Organization. *The ICD-10, Classification of mental and behavioural disorders*, Geneva: WHO, 1994.

4. Cardeña E, Nijenhuis E. Embodied sorrow: a special issue on somatoform dissociation. *J Trauma Dissociation* 2000; 1: 1-6

5. Lipsanen T, Saarijärvi S, Lauerma H. Exploring the relations between depression, somatization, dissociation, and alexithymia - overlapping or independent constructs? *Psychopathology* 2004; 37: 200-206

6. Nemiah JC. Dissociation, conversion, and somatization. In: Spiegel D, ed. *Dissociative disorders: A clinical review*. Lutherville: Sidran Press, 1993: 104-117

7. Saxe GN, Chinman G, Berkowitz R, Hall K, Lieberg G, Schwartz J, Van der Kolk BA. Somatization in patients with dissociative disorders. *Am J Psychiatry* 1994; 151: 1329-1334

8. Spitzer C, Spelsberg B, Grabe H-J, Mundt B, Freyberger HJ. Dissociative experiences and psychopathology in conversion disorders. *J Psychosom Res* 1999; 46: 291-294

- 1
2
3
4
5 9. Boon S, Draijer N. Diagnosing dissociative disorders in the Netherlands: a pilot study
6
7 with the structured interview for DSM-III-R dissociative disorders. *Am J Psychiatry*
8
9 1991; 148: 458-462
10
- 11 10. Coons PM. Dissociative disorder not otherwise specified: a clinical investigation of
12
13 50 cases with suggestions for typology and treatment. *Dissociation* 1992; 5: 187-195
14
- 15 11. Coons PM. Psychogenic amnesia: a clinical investigation of 25 cases. *Dissociation*
16
17 1992; 5: 73-79
18
- 19 12. Nijenhuis ERS, Spinhoven P, Van Dyck R, Van der Hart O, Vanderlinden J.
20
21 Psychometric characteristics of the somatoform dissociation questionnaire: a replication
22
23 study. *Psychother Psychosom* 1998; 67: 17-23
24
- 25 13. Nijenhuis ERS, Van Dyck R, Spinhoven P, *et al.* Somatoform dissociation
26
27 discriminates among diagnostic categories over and above general psychopathology.
28
29 *Aust N Z J Psychiatry* 1999; 33: 511-520
30
- 31 14. Ross CA, Fast E, Anderson G, Auty A, Todd J. Somatic symptoms in multiple
32
33 sclerosis and MPD. *Dissociation* 1990; 3: 102-106
34
- 35 15. Ross CA, Miller SD, Reagor P, Bjornson L, Fraser GA, Anderson G. Structured
36
37 interview data on 102 cases of multiple personality disorder from four Centers. *Am J*
38
39 *Psychiatry* 1990; 147: 596-601
40
- 41 16. Sar V, Kundakçı T, Kızıltan E, Bakim B, Bozkurt O. Differentiating dissociative
42
43 disorders from other diagnostic groups through somatoform dissociation in Turkey. *J*
44
45 *Trauma Dissociation* 2000; 1: 67-80
46
- 47 17. Walker NG, Katon WJ, Neraas K, Jemelka RP, Massoth D. Dissociation in women
48
49 with chronic pelvic pain. *Am J Psychiatry* 1992; 149: 534-537
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 18. Martínez-Taboas A. Multiple personality disorder in Puerto Rico: analysis of fifteen
5 cases. *Dissociation* 1991; 4: 189-192
6
7
8
9 19. Brown RJ, Schrag A, Trimble MR. Dissociation, childhood interpersonal trauma,
10 and family functioning in patients with somatization disorder. *Am J Psychiatry* 2005;
11 162: 899-905
12
13
14 20. Pribor EF, Yutzy SH, Dean JT, Wetzel RD. Briquet's syndrome, dissociation, and
15 abuse. *Am J Psychiatry* 1993; 150: 1507-1511
16
17
18 21. Guz H, Doganay Z, Ozkan A, Colak E, Tomac A, Sarisoy G. Conversion and
19 somatization disorders: the dissociative symptoms and other characteristics. *J*
20 *Psychosom Res* 2004; 56: 287-291
21
22
23 22. Spitzer C, Freyberger HJ, Stieglitz R-D, *et al.* Adaptation and psychometric
24 properties of the German version of the dissociative experiences scale. *J Trauma Stress*
25 1998; 11: 799-809
26
27
28 23. Tezcan E, Atmaca M, Kuloglu M, Gecici O, Buyukbayram A, Tutkun H.
29 Dissociative disorders in Turkish inpatients with conversion disorder. *Compr Psychiatry*
30 2003; 44: 324-330
31
32
33 24. Akyüz G, Kugu N, Akyüz A, Dogan O. Dissociation and childhood abuse history in
34 epileptic and pseudoseizure patients. *Epileptic Disord* 2006; 6: 187-192
35
36
37 25. Bowman ES, Markand ON. Psychodynamics and psychiatric diagnoses of
38 pseudoseizure subjects. *Am J Psychiatry* 1996; 153: 57-63
39
40
41 26. D' Alessio L, Giagante B, Oddo S, *et al.* Psychiatric disorders in patients with
42 psychogenic non-epileptic seizures, with and without comorbid epilepsy. *Seizure* 2006;
43 15: 333-339
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

27. Harden CL. Pseudoseizures and dissociative disorders: A common mechanism involving traumatic experiences. *Seizure* 1997; 6: 151-155
28. Prueter C, Schultz-Venrath U, Rimpau W. Dissociative and associated psychopathological symptoms in patients with epilepsy, pseudoseizures, and both seizure forms. *Epilepsia* 2002; 43: 188-192
29. Van Merode T, Twellaar M, Kotsopoulos IAW, *et al.* Psychological characteristics of patients with newly developed psychogenic seizures. *J Neurol Neurosurg Psychiatry* 2004; 75: 1175-1177
30. Bowman ES. Why conversion seizures should be classified as a dissociative disorder. *Psychiatr Clin North Am* 2006; 29: 185-211
31. Cardeña E, Spiegel D. Diagnostic issues, criteria and comorbidity of dissociative disorders. In: Michelson LK, Ray WJ, eds. *Handbook of dissociation: theoretical, empirical, and clinical perspectives*. New York: Plenum, 1996: 227-250.
32. Janet P. *The mental state of hystericals*. New York: Putnam, 1901.
33. Kent DA, Tomasson K, Coryell W. Course and outcome of conversion and somatization disorders: a four- year follow-up. *Psychosomatics*, 1995; 36: 138-144
34. Näring G, Nijenhuis ERS. Relationships between self-reported potentially traumatizing events, psychoform and somatoform dissociation, and absorption, in two non-clinical populations. *Aust N Z J Psychiatry* 2005; 39: 982-988
36. Nijenhuis ERS, Spinhoven P, Van Dyck R, Van der Hart O, Vanderlinden J. The development and psychometric characteristics of the somatoform dissociation questionnaire (SDQ-20). *J Nerv Ment Dis* 1996; 184: 688-694

37. Van der Hart O, Van Dijke A, Van Son M, Steele K. Somatoform dissociation in traumatized World War I combat soldiers: A neglected clinical heritage. *J Trauma Dissociation* 2000; 1: 33-65
38. Waller G, Hamilton K, Elliot P, *et al.* Somatoform dissociation, psychological dissociation, and specific forms of trauma. *J Trauma Dissociation* 2000; 1: 81-98
39. Bernstein EM, Putnam FW. Development, reliability and validity of a dissociation scale. *J Nerv Ment Dis* 1986; 174: 727-735
40. Spitzer RL. Psychiatric diagnosis: are clinicians still necessary? *Compr Psychiatry* 1983; 24: 399-411
41. Ross CA, Heber S, Norton GR, Anderson G, Anderson D, Barchet P. The dissociative disorders interview schedule: a structured interview. *Dissociation* 1989; 2: 169-189
42. Espirito Santo H, Pio Abreu JL. Portuguese validation of the dissociative experiences scale (DES). *J Trauma Dissociation* in press
43. Espirito Santo H, Pio Abreu JL. Dissociative disorders and other psychopathological groups: exploring the differences through somatoform dissociation questionnaire (SDQ-20). *Rev Bras Psiquiatr* 2007; 29: 354-358
44. Derogatis LR, Melisaratos N. The brief symptom inventory: an introductory report. *Psychol Med* 1983; 13: 596-605
45. Canavarro MC. Inventário de sintomas psicopatológicos. In: Simões MR, Gonçalves MM, Almeida LS, eds. *Testes e provas projectivas em Portugal*, vol. 2. Braga: APPORT, 1988: 95-109.

46. Espirito Santo HA, Madeira F, Pio Abreu JL. Versão portuguesa do dissociative disorders interview schedule (DDIS-P), Estudo preliminar de adaptação a uma amostra da população portuguesa. *Rev Port Psiquiatr* 2007; 28: 5-17
47. Kendell R, Jablensky A. Distinguishing between the validity and utility of psychiatric diagnoses. *Am J Psychiatry* 2003; 160: 4-12
48. Wessely S, Nimnuan C, Sharpe M. Functional somatic syndromes: one or many? *Lancet* 1999; 354: 936-939
49. Mayou RA. Medically unexplained physical symptoms. *Br Med J* 1991; 303: 534-535.
50. Krüger C, Van Staden W. Is conversion a dissociative symptom? *Bridging Eastern and Western Psychiatry* 2003; 1: 88-94
51. Sar V, Akyüz G, Kundakçi T, Kiziltan E, Dogan O. Childhood trauma, dissociation, and psychiatric comorbidity in patients with conversion disorder. *Am J Psychiatry* 2004; 161: 2271-2276
52. Sharpe M, Mayou R. Somatoform disorders: a help or hindrance to good patient care? *Br J Med Psychol* 2004; 184: 465-467
53. Moene FC, Spinhoven P, Hoogduin K, Sandyck P, Roelofs K. Hypnotizability, dissociation and trauma in patients with a conversion disorder: an exploratory study. *Clin Psychol Psychother* 2001; 8: 400-410
54. World Health Organization. *Gender and women's mental health*. Geneva: WHO, 2006. [cited 21 Dec 2006.] Available from http://www.who.int/mental_health/prevention/genderwomen/en/print.html

- 1
2
3
4
5 55. Dorahy MJ, Lewis CA, Millar RG, Gee TL. Predictors of nonpathological
6
7 dissociation in Northern Ireland: the affects of trauma and exposure to political
8
9 violence. *J Trauma Stress* 2003; 16: 611-615
10
11
12 56. Espirito Santo HMA, Pio-Abreu JL. Demographic and mental health factors
13
14 associated with pathological dissociation in a Portuguese sample. *J Trauma*
15
16 *Dissociation* 2008; 9: 369-387
17
18
19 57. Maaranen P, Tanskanen A, Kaisa H, *et al.* The relationship between psychological
20
21 and somatoform dissociation in the general population. *J Nerv Ment Dis* 2005; 193:
22
23 690-692
24
25
26 58. Spitzer C, Barnow S, Grabe H-J, *et al.* Frequency, clinical and demographic
27
28 correlates of pathological dissociation in Europe. *J Trauma Dissociation* 2006; 7: 51-56
29
30
31 59. Seedat S, Stein MB, Forde DR. Prevalence of dissociative experiences in a
32
33 community sample: relationship to gender, ethnicity, and substance use. *The J Nerv*
34
35 *Ment Dis* 2003; 191: 115-120
36
37
38 60. Maaranen P, Tanskanen A, Haatainen K, Koivumaa-Honkanen H, Hintikka J,
39
40 Viinamäki H. Somatoform dissociation and adverse childhood experiences in the
41
42 general population. *J Nerv Ment Dis* 2004; 192: 337-342
43
44
45 61. Saxe GN, Van der Kolk BA, Berkowitz R, *et al.* Dissociative disorders in
46
47 psychiatric inpatients. *Am J Psychiatry* 1993; 150: 1037-1042
48
49
50 62. Stone J, Sharpe M, Binzer M. Motor conversion symptoms and pseudoseizures: A
51
52 comparison of clinical characteristics. *Psychosomatics* 2004; 45: 492-499
53
54
55
56
57
58
59
60

Table 1. Demographic characteristics of psychopathological groups.

Groups	Total	Age		Gender		Marital Status		School	
		(years)		(M/F)		(married/single)		Years	
	<i>n</i>	<i>M</i>	<i>SD</i>	<i>n</i>	%	<i>n</i>	%	<i>M</i>	<i>SD</i>
Dissociative Disorders	39	34.0	11.7	11/28	28.2/71.8	20/19	51.3/48.7	9.7	4.5
Conversion Disorder	26	27.4	8.8	6/20	23.1/76.9	8/18	30.8/69.2	10.9	4.1
Somatization Disorder	40	32.6	12.8	14/26	35.0/65.0	15/25	37.5/62.5	11.4	3.7
Other Psychiatric Disorders	46	31.5	11.3	14/32	30.4/69.6	18/28	39.1/60.9	11.7	4.4
Total	151	31.7	11.5	45/106	29.8/70.2	61/90	40.4/59.6	10.9	4.2

Table 2. Frequencies of group pathologies and their main disorder frequencies (N=151).

Group pathologies	Disorders	n	%
Conversion disorders		26	17.2
	Pseudo seizure	1	0.7
	Combination	3	2.0
	Motor	10	6.6
	Sensorial	12	7.9
Dissociative Disorders		39	25.8
	DDOS	10	6.6
	Fugue	7	5.3
	Amnesia	11	7.3
	Depersonalization	11	7.3
Somatization disorder		40	26.5
Other psychiatric disorders		46	30.5
	Panic disorder	8	5.3
	Specific Phobias	8	5.3
	Social Phobia	9	6.0
	Depression	10	6.6
	OCD	11	7.3

Table 3. Psychoform dissociation (DES), somatoform dissociation (SDQ), global severity index (GSI) and psychopathological symptoms in patients with conversion disorder, dissociative disorders, somatization disorder, and the control group (N=151).

Measures	Conversion disorder (CV) (n=26)		Dissociative Disorders (DD) (n=39)		Somatization Disorder (So) (n=40)		Other psychiatric disorders (OP) (n=46)		Kruskal-Wallis (df=3)	
	M	SD	M	SD	M	SD	M	SD	H	p
DES	43.49	12.33	36.02	10.10	19.39	11.89	18.02	8.37	68.86	0.000
SDQ	39.76	14.15	39.28	11.88	31.81	9.16	29.61	7.06	20.46	0.000
BSI										
GSI	1.48	0.43	1.55	0.57	1.54	0.54	1.23	0.43	10.40	0.015
Somatization	2.00	0.92	1.05	0.78	1.74	0.75	0.89	0.59	27.30	0.000
Interpersonal sensitivity	1.46	0.69	1.42	0.82	1.63	0.72	1.43	0.66	1.39	0.709
Anxiety	1.71	0.82	1.39	0.65	1.54	0.80	1.30	0.64	3.40	0.335
Phobic Anxiety	0.97	0.97	0.52	0.65	0.85	0.83	0.92	0.73	4.30	0.232
Psychoticism	1.43	0.93	1.53	0.93	1.17	0.65	1.10	0.63	4.92	0.178
Obsessive-compulsion	1.71	0.67	1.70	0.80	1.74	0.66	1.48	0.59	2.53	0.471
Depression	1.50	0.85	1.98	1.20	1.56	0.86	1.50	0.73	1.60	0.659
Hostility	1.06	0.68	0.97	0.78	1.32	0.87	1.09	0.67	1.52	0.679
Paranoid ideation	1.69	0.69	1.78	0.77	1.59	0.68	1.37	0.58	3.95	0.267
Pairwise Post-hoc Mann-Whitney										
DES: CV>So ***; CV>OP**; DD>So***; DD>OP***										
Other comparisons non significant.										
SDQ: CV>OP***; DD>So*; DD>OP*										
Other comparisons non significant.										
GSI: comparisons non significant.										
Somatization: Co>OP***, So>OP***										

* p<0.0083, ** p<0.0017, *** p<0.00017 (Bonferroni-corrected)

Table 4. Percentages of cases of diagnostic categories scoring above the cut-offs on the Dissociative Experiences Scale (DES) and the Somatoform Dissociation Questionnaire (SDQ-20) (N= 151).

Pathologies	DES≥30		SDQ≥35	
	n	%	n	%
Conversion disorders (CV)	29	78.4	22	59.5
Dissociative disorders (DD)	18	69.2	15	60.0
Somatization disorder (So)	6	15.0	10	25.0
Other psychiatric disorders (OP)	5	10.9	10	21.7
Kruskal-Wallis (df=3)	58.73***		20.14***	
Pairwise Post-hoc Mann-Whitney				
DES: CV>So ***; CV>OP***; DD>So***; DD>OP***				
Other comparisons non significant.				
SDQ: CV>So*; CV>OP**; DD>So*; DD>OP***				
Other comparisons non significant.				
* p<0.0083, ** p<0.0017, *** p <0.00017 (Bonferroni-corrected)				